

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/143052>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.

*Journal of Organometallic Chemistry*, 65 (1974) C43—C45  
© Elsevier Sequoia S.A., Lausanne — Printed in The Netherlands

# Preliminary communication

## SOME ADDUCTS OF $\text{RhCl}[\text{P}(\text{C}_6\text{H}_{11})_3]_2$ , A RHODIUM COMPOUND SPONTANEOUSLY COORDINATING DINITROGEN

H.L.M. VAN GAAL, F.G. MOERS and J.J. STEGGERDA

*Department of Inorganic Chemistry, Catholic University, Toernooiveld, Nijmegen  
(The Netherlands)*

(Received October 25th, 1973)

### Summary

Solutions of  $\text{RhCl}(\text{PCy}_3)_2$  react with  $\text{O}_2$ ,  $\text{N}_2$ ,  $\text{C}_2\text{H}_4$ ,  $\text{CO}$  and  $\text{H}_2$  to give *trans*- $\text{RhClA}(\text{PCy}_3)_2$  ( $\text{A} = \text{O}_2$ ,  $\text{N}_2$ ,  $\text{C}_2\text{H}_4$  and  $\text{CO}$ ,  $\text{Cy} = \text{cyclohexyl}$ ) and  $\text{RhClH}_2(\text{PCy}_3)_2$ ; the spontaneous formation of the rather air-stable  $\text{RhCl}(\text{PCy}_3)_2\text{N}_2$  is ascribed to a combination of the steric requirements and electronic properties of the phosphine ligand.

Tricyclohexylphosphine ( $\text{PCy}_3$ ), like other bulky phosphines, is effective in stabilizing unusual coordination numbers and valence states [1, 2]. In the reaction of  $\text{PCy}_3$  with  $\text{RhCl}_3 \cdot 3\text{H}_2\text{O}$  no reduction to the  $\text{Rh}^{\text{I}}$  valence state occurs, but instead *inter alia*  $\text{Rh}^{\text{III}}\text{Cl}_2(\text{PCy}_3)_2$  is formed [2]. Although the existence of  $\text{Rh}^{\text{III}}\text{Cl}_3(\text{PCy}_3)_3$  has been reported [3], we did not succeed in its preparation and we have not found an indication for a  $\text{PCy}_3/\text{Rh}$  ratio exceeding 2 in any other complex. In this paper we report the preparation of  $\text{Rh}^{\text{I}}$  tricyclohexylphosphine compounds by cyclooctene displacement from  $[\text{RhCl}(\text{C}_8\text{H}_{14})_2]_2$  [4].

Addition of two moles of  $\text{PCy}_3$  per mole of  $[\text{RhCl}(\text{C}_8\text{H}_{14})_2]_2$  resulted in the rapid formation of ochrous  $[\text{RhCl}(\text{PCy}_3)(\text{C}_8\text{H}_{14})]_2$  (I). The solvated  $\text{C}_8\text{H}_{14}$ -free complex  $\text{RhCl}(\text{PCy}_3)_2(\text{S})$  ( $\text{S} = \text{solvent molecule}$ ) was rather slowly formed (20 min stirring) by reaction of either 4 moles or an excess (8 moles) of  $\text{PCy}_3$  ( $\text{Rh}/\text{PCy}_3 = 1/2$ , resp.  $1/4$ ) with  $[\text{RhCl}(\text{C}_8\text{H}_{14})_2]_2$  in  $\text{C}_6\text{H}_6$ , and was isolated as a lilac precipitate of yet not fully identified nature. When  $[\text{RhCl}(\text{C}_2\text{H}_4)_2]_2$  was used, yellow *trans*- $\text{RhCl}(\text{C}_2\text{H}_4)(\text{PCy}_3)_2$  (II) was obtained. The lilac product is unstable in the solid state, it decomposes within a few hours under vacuum and somewhat more slowly under nitrogen. It immediately turns brown upon exposure to air. In solution, however,  $\text{RhCl}(\text{PCy}_3)_2(\text{S})$  reacts rapidly with oxygen to give the grayish-blue adduct *trans*- $\text{RhCl}(\text{O}_2)(\text{PCy}_3)_2$  (III), while the dinitrogen adduct *trans*- $\text{RhCl}(\text{N}_2)(\text{PCy}_3)_2$  (IV) is formed when a solution of

$\text{RhCl}(\text{PCy}_3)_2(\text{S})$  is exposed to 1 atm of  $\text{N}_2$  for five days. CO equally adds to  $\text{RhCl}(\text{PCy}_3)_2(\text{S})$  to yield the known *trans*- $\text{RhCl}(\text{CO})(\text{PCy}_3)_2$  (V) [2].

$\text{RhCl}(\text{PCy}_3)_2(\text{S})$ , made in situ, catalyses the hydrogenation of cyclohexene in benzene at a rate which is about 40 times less than the rate of hydrogenation by  $\text{RhCl}(\text{PPh}_3)_3$  under similar conditions [5]. The yellow dihydride  $\text{RhClH}_2(\text{PCy}_3)_2$  (VI) and the corresponding  $\text{RhClD}_2(\text{PCy}_3)_2$  (VII) can most conveniently be prepared by the reaction of  $\text{RhCl}(\text{PCy}_3)_2(\text{C}_2\text{H}_4)$  with  $\text{H}_2$  or  $\text{D}_2$ .

Satisfactory analyses of the compounds have been obtained. Infrared spectroscopic data are given in Table 1. The large variation in Rh—Cl stretching vibration frequencies suggests that in the complexes  $\text{RhClA}(\text{PCy}_3)_2$ , Cl and A are in *trans* positions as are the two bulky  $\text{PCy}_3$  ligands. The dihydride (VI) has a different structure and its Rh—Cl stretching vibration frequency may not be compared with those of the other adducts.

TABLE 1

INFRARED SPECTROSCOPIC DATA FOR COMPOUNDS  $\text{RhCl}(\text{PCy}_3)_2\text{A}$ 

A	IR absorptions ( $\text{cm}^{-1}$ )	
	$\nu(\text{Rh—Cl})$	others
$\text{C}_2\text{H}_4$	294m	3077w, 3040w, 3017w ( $\nu(\text{CH})$ ); 1510w (br), 1208m, 1183w; 950m, 933w ( $\delta(\text{CH})$ )
CO	304m	1942vs ( $\nu(\text{CO})$ ), 584s ( $\delta(\text{CO})$ )
$\text{N}_2$	317m	2103vs ( $\nu(\text{NN})$ ), 470m ( $\nu(\text{RhN})$ )
$\text{O}_2$	328m	993m <sup>a</sup>
$\text{H}_2$	291m	2165(sh), 2120m ( $\nu(\text{Rh—H})$ ), 622m(br) ( $\delta(\text{Rh—H})$ )
$\text{D}_2$	291m	1560(sh), 1528m ( $\nu(\text{Rh—D})$ )

<sup>a</sup>Not assigned.  $\text{PCy}_3$  absorbs in the 800-900  $\text{cm}^{-1}$  region where  $\text{MO}_2$  modes normally are found [7].

We assume the bulkiness of tricyclohexylphosphine to be one of the causes of the formation of the dinitrogen adduct (IV). For  $\text{PCy}_3$ , neither formation of a tris-complex comparable to  $\text{RhCl}(\text{PPh}_3)_3$  [5] nor of a dimer comparable to  $[\text{RhCl}(\text{PPh}_3)_2]_2$  [5] seems possible. Also adducts with  $\text{C}_6\text{H}_6$  or  $\text{C}_8\text{H}_{14}$ , the other molecules present, seem to be labilized by steric influences of the two  $\text{PCy}_3$  ligands. We therefore postulate the transient existence of the 14 metal valence electron species  $\text{RhCl}(\text{PCy}_3)_2$  as the  $\text{N}_2$ -bonding species to account for this spontaneous dinitrogen coordination, which has not been reported before for rhodium(I) complexes. The comparable rhodium—dinitrogen compound  $\text{RhCl}(\text{PPh}_3)_2(\text{N}_2)$  has been prepared by an indirect method [6]. The high basicity of  $\text{PCy}_3$  may facilitate the addition of  $\text{N}_2$ , and certainly improves the stability of  $\text{RhCl}(\text{PCy}_3)_2(\text{N}_2)$ , which is only partly decomposed by air upon standing overnight in  $\text{C}_6\text{H}_6$  or  $\text{CHCl}_3$ . At room temperature the  $\text{N}_2$ -ligand can be replaced by CO, but not by  $\text{C}_2\text{H}_4$  or  $\text{H}_2$ . In similar reactions CO displaces  $\text{C}_2\text{H}_4$ ,  $\text{H}_2$ , and  $\text{O}_2$  from their adducts, to yield spectroscopically pure (V). Oxygen slowly replaces  $\text{C}_2\text{H}_4$  and  $\text{H}_2$  to give impure (III). In contrast to the behaviour of  $\text{RhCl}(\text{PPh}_3)_2\text{A}$  (A =  $\text{C}_2\text{H}_4$  or  $\text{H}_2$ ) [5],  $\text{RhCl}(\text{PCy}_3)_2\text{A}$  appears not to lose its coordinated molecule of  $\text{C}_2\text{H}_4$  or  $\text{H}_2$  on sweeping its solution with nitrogen.

A mixture of  $[\text{IrCl}(\text{C}_8\text{H}_{14})_2]_2$  [4] and  $\text{PCy}_3$  in  $\text{C}_6\text{H}_6$  does not coordinate dinitrogen. Reactions with this system and further reactions with the rhodium system are currently under investigation.

We thank Mr. J. Diersmann for the elemental analyses and Mrs. A.W.P.G. Peters-Rit for the kinetic measurements.

## References

- 1 F.G. Moers and J.P. Langhout, *Rec. Trav. Chim. Pays-Bas*, 91 (1972) 591.
- 2 F.G. Moers, J.A.M. de Jong and P.M.H. Beaumont, *J. Inorg. Nucl. Chem.*, 35 (1973) 1915.
- 3 G.M. Intille, *Inorg. Chem.*, 11 (1972) 695.
- 4 A. van der Ent and A.L. Onderdelinden, *Inorg. Synth.*, 14 (1973) 92.
- 5 J.A. Osborn, F.H. Jardine, J.F. Young and G. Wilkinson, *J. Chem. Soc. A*, (1966) 1711.
- 6 L.Yu. Ukhin, Yu.A. Shvetsov and M.L. Khidekel', *Bull. Acad. Sci. U.S.S.R., Div. Chem. Sci.*, (1967) 934.
- 7 V.J. Choy and C.J. O'Connor, *Coord. Chem. Rev.*, 9 (1972) 145.

